

## 2018年度日本政府(文部科学省)奨学金留学生選考試験

## QUALIFYING EXAMINATION FOR APPLICANTS FOR JAPANESE GOVERNMENT (MEXT) SCHOLARSHIPS 2018

学科試験 問題 EXAMINATION QUESTIONS

(学部留学生) UNDERGRADUATE STUDENTS

> 数 学(A) MATHEMATICS(A)

注意 ☆試験時間は60分。 PLEASE NOTE: THE TEST PERIOD IS **60 MINUTES**.



## MATHEMATICS(A)

(2018)

				] .	
Nationality		No.			Manka
Name	(Please print full name, underlining family name)			Marks	

- 1. Answer the following questions and fill in your responses in the corresponding boxes on the answer sheet.
  - (1) The number of digits of  $7^{2677}$  is [1-1] and the last digit of it is [1-2], where  $\log_{10} 3 = 0.4771$ ,  $\log_{10} 7 = 0.8451$ .
  - (2) Simplify  $\frac{1}{1+\sqrt{2}} + \frac{1}{\sqrt{2}+\sqrt{3}} + \frac{1}{\sqrt{3}+\sqrt{4}} + \frac{1}{\sqrt{4}+\sqrt{5}}$  as briefly as possible. The result is [1-3].
  - (3) Assume that  $0 < \theta < \frac{\pi}{4}$ . If  $\sin 2\theta = \frac{1}{4}$ , then  $\frac{\sin \theta + \cos \theta}{-\sin \theta + \cos \theta} = \boxed{[1-4]}$
  - (4) Let  $P_1, P_2, P_3, P_4, P_5$ , and  $P_6$  be the vertices of a regular hexagon in anticlockwise order. We throw a fair dice three times and denote the scores shown on the dice as the ordered triple (i, j, k). In this case, the probability that the three points  $P_i, P_j, P_k$  make a triangle is 1-5.
  - (5) For the equation  $4^x 2^x 12 = 0$ , the real solution is  $x = \boxed{1-6}$ .
  - (6) For a pyramid OABC, the centroids of the triangles OAB, OBC, and OCA are F, G, and H, respectively. For the centroid P of the triangle FGH, the vector  $\overrightarrow{OP}$  is given by

$$\vec{OP} \ = \ \frac{2}{\boxed{\boxed{[1\text{-}7]}}} \left( \vec{OA} + \vec{OB} + \vec{OC} \right).$$



- (7) Let a point O be the origin of xy-coordinate plane. We define four points A(1, 0), B(1, 1), C(2, 1), D(3, 1) on the plane. Let us start from C, go through a point on line OA, go through a point on line OB, and reach D with the minimum length of the path. In this path, the point on the line OA is ([1-8], [1-9]), that on the line OB is ([1-10], [1-11]), the length of the path is [1-12].
- (8) Assume that integers m and n satisfy  $2|m|+3|n-1| \le 7$ . m+n is maximum when  $(m, n) = \left(3, \boxed{[1\text{-}13]}\right)$ ,  $\left(\boxed{[1\text{-}14]}, \boxed{[1\text{-}15]}\right)$  and its maximum value is  $\boxed{[1\text{-}16]}$ .
- (9) If a quadratic function f(x) is maximum at x = 1 with the maximum value 5, and satisfies f(-2) = -22, it is given by

$$f(x) = [1-17]x^2 + [1-18]x + [1-19].$$

(10) When integers k and n satisfy  $1 \le k \le n$ , we have

$$\sum_{l=k}^{n} 2^{l} = 2^{\left[1-20\right]} - 2^{\left[1-21\right]}.$$

Therefore, it follows that

$$\sum_{k=1}^{n} k 2^{k} = \sum_{k=1}^{n} \sum_{l=k}^{n} 2^{l} = \left( \boxed{[1-22]} \right) 2^{\boxed{[1-23]}} + 2.$$

(11) A decimal number 123456 is shown by a ternary (base 3) number [1-24].

(Describe only the value of the ternary number without describing a notation that indicates a ternary numeral system.)



- **2.** For a cubic function  $f(x) = x^3 3ax^2 + 3bx 2$ , answer the following questions and fill in your responses in the corresponding boxes on the answer sheet.
  - (1) If x = 1, 3 are the extreme points of f(x), then a = [2-1] and b = [2-2]. In this case, the solutions of f(x) = 0 can be arranged as [2-3] < [2-4] < [2-5] in increasing order.
  - (2) Assume that a = b. If the function f(x) is monotonously increasing, then  $[2-6] \le a \le [2-7]$ .



**3.** In xyz-coordinate system, we define a solid A by

$$\frac{1}{9}x^2 + \frac{1}{4}y^2 \le z^4 \quad (0 \le z \le 1).$$

Fill in your responses in the corresponding boxes on the answer sheet.

(1) We define a solid B by

$$x^2 + y^2 \le z^4 \quad (0 \le z \le 1)$$

The volume of the solid B is [3-1].

- (2) The solid A is given by elongating the solid B [3-2] times in the x-axis direction and [3-3] times in the y-axis direction.
- (3) The volume of the solid A is [3-4] times as large as that of the solid B.
- (4) The volume of the solid A is [3-5].